

*The first and only IV agent for the treatment of pain and fever*

**CALDOLOR<sup>®</sup>**  
*(ibuprofen) Injection*

## From OR to ER, the IV agent for pain and fever

### **Pain reduction superior to PCA-delivered narcotics alone**

Significantly less patient-reported pain at rest and with movement following abdominal hysterectomy and orthopedic surgeries<sup>1,2</sup>

### **Significant opioid-sparing effect vs PCA-delivered narcotics**

Within first 24 hours postsurgery<sup>1,2</sup>

### **For the first time, IV temperature reduction within 30 minutes**

Significant temperature reduction after one dose<sup>1,2</sup>

## Indications and usage

CALDOLOR<sup>®</sup> is indicated in adults for the<sup>2</sup>

- Management of mild to moderate pain
- Management of moderate to severe pain as an adjunct to opioid analgesics
- Reduction of fever

## The benefits of IV

- Consistent, reliable delivery
- Provides an alternate route of administration

## The level of safety that ibuprofen delivers

- In clinical trials of over 900 patients, CALDOLOR had an adverse event profile comparable to placebo<sup>1</sup>
- No significant differences in adverse events were seen between CALDOLOR and placebo that would suggest renal, cardiac, or bleeding complications often associated with long-term administration of NSAIDs<sup>1</sup>

Please see Important Safety Information and Warning on reverse side.

The first and only IV agent for the treatment of pain and fever

# CALDOLOR<sup>®</sup> (ibuprofen) Injection

## Dosing

### Pain

Adults: 400 mg to 800 mg over 30 minutes every 6 hours as needed<sup>2</sup>

### Fever

Adults: 400-mg initial dose, over 30 minutes<sup>2</sup>

Follow with 400 mg every 4 to 6 hours or 100 mg to 200 mg every 4 hours as necessary<sup>2</sup>

Not to exceed 3200-mg daily dose<sup>2</sup>

CALDOLOR<sup>®</sup> must be diluted prior to intravenous infusion. Diluted infusions are stable for at least 7 days.



Product Specifications	400-mg/4-mL Vial (Tray of 25)	800-mg/8-mL Vial (Tray of 25)
NDC Number	66220-247-04	66220-287-08
UPC Code	[01]30366220247047	[01]30366220287081
WAC	\$183.75	\$262.50
AWP	\$230.00	\$328.15
Expiration Date	5 years	5 years
Dimensions	4.11" x 1.8" x 4.11"	4.5" x 1.5" x 5.59"

### WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

#### Cardiovascular Risk

- Nonsteroidal anti-inflammatory drugs (NSAIDs) may increase the risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. Risk may increase with duration of use
- CALDOLOR is contraindicated for the treatment of perioperative pain in the setting of coronary artery bypass graft (CABG) surgery

#### Gastrointestinal Risk

- NSAIDs increase the risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. Events can occur at any time without warning symptoms. Elderly patients are at greater risk

### IMPORTANT SAFETY INFORMATION

CALDOLOR should be used with caution in patients with congestive heart failure, kidney impairment, at risk of blood clots, and in those who have a history of ulcers or gastrointestinal bleeding. When used in such patients, attention to using the lowest effective dose for the shortest time period is important to reduce the risk of serious adverse events. Ibuprofen has been associated with high blood pressure, serious skin reactions, and serious allergic reactions.

The most common adverse events reported in the controlled clinical trials were nausea, flatulence, vomiting, and headache.

Please see full Prescribing Information, including Boxed Warning.

CALDOLOR is a registered trademark of Cumberland Pharmaceuticals Inc.

**References:** 1. Data on file, Cumberland Pharmaceuticals Inc., Nashville, TN. 2. Caldolor [prescribing information]. Nashville, TN: Cumberland Pharmaceuticals Inc; 2009.

**Contact your wholesaler to order CALDOLOR.**

For medical inquiries, contact Professional Affairs: 1-877-484-2700  
www.CALDOLOR.com

